

THE SERUM VITAMIN D STATUS IN ADULTS WITH DIABETIC RETINOPATHY: A SYSTEMATIC REVIEW AND META-ANALYSIS

STATUS VITAMINA D U SERUMU KOD ODRASLIH SA DIJABETIČKOM RETINOPATIJOM: SISTEMATSKI PREGLED I META-ANALIZA

Hongyu Wei^{1,2}, Hao Wang², Ke Diao², Xiaoxuan Wang²,
Hong Chen², Congying Wang², Minglian Zhang^{1,2*}

¹Graduate School of Hebei University of Chinese Medicine, Shijiazhuang, China

²Hebei Eye Hospital, The Affiliated Hospital of Hebei University of Chinese Medicine Xingtai, China

Summary

Background: Emerging evidence suggests that vitamin D may play a crucial role in the development of diabetic retinopathy (DR). However, the extent of vitamin D deficiency in individuals with DR remains uncertain. This study aimed to evaluate serum vitamin D levels in adults with DR.

Methods: A literature was retrieved using the Embase, PubMed and Cochrane databases to identify observational studies that evaluated the levels of 25(OH)D₃, 1,25(OH)₂D₃ or total 25(OH)D in the serum of patients with DR. A total of 31 independent studies were included for meta-analysis.

Results: The pooled mean concentration of total 25(OH)D among individuals with diabetes was 16.04 ng/mL (95% CI: 15.13–16.96; I² = 98.8%), while the mean serum level of 25(OH)D₃ in those with DR was 10.68 ng/mL (95% CI: 5.94–15.41; I² = 99.5%), both significantly below the normal reference range. The average concentration of 1,25(OH)₂D₃ was 31.14 pg/mL (95% CI: 24.35–37.94; I² = 98.8%).

Conclusions: The evidence from this meta-analysis indicates an association between vitamin D deficiency and an increased risk of DR.

Keywords: diabetic retinopathy, vitamin D, 25-hydroxy-vitamin D₃, 1,25-dihydroxyvitamin D₃, meta-analysis

Kratak sadržaj

Uvod: Novi dokazi ukazuju na to da vitamin D može igrati ključnu ulogu u razvoju dijabetičke retinopatije (DR). Međutim, stepen nedostatka vitamina D kod osoba sa DR ostaje neizvestan. Cilj ove studije bio je da se procene nivoi vitamina D u serumu kod odraslih sa DR.

Metode: Prikupljena je literatura korišćenjem baza podataka Embase, PubMed i Cochrane kako bi se identifikovale opservacione studije koje su procenjivale nivoe 25(OH)D₃, 1,25(OH)₂D₃ ili ukupnog 25(OH)D u serumu pacijenata sa DR. Ukupno 31 nezavisna studija je uključena u meta-analizu.

Rezultati: Zbirna prosečna koncentracija ukupnog 25(OH)D kod osoba sa dijabetesom bila je 16,04 ng/mL (95% CI: 15,13–16,96; I² = 98,8%), dok je prosečan nivo 25(OH)D₃ u serumu kod osoba sa DR bio 10,68 ng/mL (95% CI: 5,94–15,41; I² = 99,5%), oba značajno ispod normalnog referentnog opsega. Prosečna koncentracija 1,25(OH)₂D₃ bila je 31,14 pg/mL (95% CI: 24,35–37,94; I² = 98,8%).

Zaključak: Dokazi iz ove meta-analize ukazuju na vezu između nedostatka vitamina D i povećanog rizika od DR.

Ključne reči: dijabetička retinopatija, vitamin D, 25-hidroksivitamin D₃, 1,25-dihidroksivitamin D₃, meta-analiza

Address for correspondence:

Minglian Zhang, BM.

Graduate School of Hebei University of Chinese Medicine,
No. 326, Xinshi South Road, Qiaoxi District, Shijiazhuang
050091, China; Hebei Eye Hospital, The Affiliated Hospital
of Hebei University of Chinese Medicine No. 399, Quanbei
East Street, Xiangdu District, Xingtai 054001, China
Tel: 86015631901999

e-mail: zhmyk@sohu.com

Introduction

DR stands as the most common ocular problems globally and serves as the predominant factor contributing to sight loss among people afflicted with diabetes (1). Statistical data indicate that roughly a third of the world's populace diagnosed with diabetes will experience DR (2). The etiology of DR is multifactorial, encompassing long-term hyperglycemia, oxidative stress, inflammatory response, vascular endothelial dysfunction and other links (3). Vitamin D, a crucial lipid-soluble vitamin and steroid hormone, is vital for regulating calcium and phosphorus metabolism and modulating immune responses in the body (4). In recent years, a growing collection of studies has centered on the influence of vitamin D in the growth and spread of DR. Research indicates that individuals with DR frequently display diminished amounts of vitamin D, implying that a decrease in vitamin D could potentially accelerate the advancement of DR (5). Vitamin D is stored in the body primarily as 25(OH)D₃. This form undergoes a two-step hydroxylation (in the liver and kidneys) to become the physiologically active form, 1,25(OH)₂D₃ (6). Serum 25(OH)D₃ levels indicate vitamin D nutritional status, while 1,25(OH)₂D₃ levels reflect its physiological activity (7). Studies show that individuals with diabetic retinopathy (DR) have significantly lower serum levels of both 25(OH)D₃ and 1,25(OH)₂D₃ compared to diabetic controls without DR (8, 9). Furthermore, vitamin D levels are inversely correlated with the severity of DR (10).

The findings from various studies exhibited a lack of complete consistency; for instance, specific studies did not identify an observable and meaningful disparity in vitamin D concentrations among the DR and untreated cohorts (11). Considering the heterogeneity and limitations of the available evidence, it is necessary to undertake a comprehensive review and quantitative assessment of pertinent studies. In this study, we systematically searched domestic and foreign databases by comprehensive evaluation, and encompassed cohort or case-control investigations that assessed serum 25(OH)D₃, 1,25(OH)₂D₃, or total 25-hydroxyvitamin D concentrations among individuals with DR, quantitatively synthesized the existing evidence, and evaluated the overall vitamin D levels of DR patients, in order to provide reference for clinical practice. At the same time, the heterogeneity and publication bias of the included studies were also assessed to judge the reliability of the results.

Materials and Methods

Methodology for searching literature

A comprehensive investigation was undertaken utilizing the Cochrane Library, PubMed, and Embase records, covering the period from the start up to September 2024. The search strategy employed

involves the integration of subject headings and free-text terms, utilizing the primary keywords comprising: »vitamin D,« »cholecalciferol,« »25-hydroxyvitamin D,« »1,25-dihydroxyvitamin D,« »diabetic retinopathy,« and »diabetes complications«. References to included studies were also retrospectively used to supplement missing studies.

Criteria for Exclusion and Inclusion

Inclusion criteria: Studies involving human subjects with DR; reporting serum 25(OH)D₃, 1,25(OH)₂D₃, or Total 25(OH)D levels; diagnosing DR using the ICO clinical grading system (12); employing case control, cohort, or cross-sectional designs. **Exclusion Criteria:** Non-original studies reviews, meetings, case reports ;animal or cell studies; duplicate publications; studies lacking extractable key data.

Data extraction and literature assessment

Two scholars conducted an independent examination of the works in question. The extracted information mainly encompassed: name of the initial writer, the year of release, the field of investigation, the sex and age of the subjects involved in the research, the description and grading of diabetes and DR, the detection method, the average and standard deviation of serum vitamin D concentrations within every cohort, along with the modification of confounding factors. Two peer reviews meticulously verified the inclusion outcomes and extracted data, engaging an additional reviewer to address any discrepancies that arose.

Quality evaluation adoption

The Newcastle-Ottawa Scale evaluates the efficacy of case-control investigations (13). The scale is scored from three aspects: choice (4 points), comparability (2 points), and exposure (3 points), aggregating to 9 points. A score of ≥ 7 is of high quality, 5–6 is of moderate quality, and < 5 is of low quality.

Data Analysis

A thorough evaluation was carried out utilizing Stata software. The effect size of each study was assessed through serum 25(OH)D₃, 1,25(OH)₂D₃, or total 25(OH), utilizing the mean of D levels along with its 95% CI. The I² test was employed to assess heterogeneity among the investigations, revealing that an I² value exceeding 50% indicated considerable variability. Once the effects were found to be heterogeneous (I² > 50%), a random effects model was used. Otherwise, a fixed effect model was used. The risk of publishing bias of the featured investigations

evaluated visually by funnel plots. To evaluate the robustness of the findings, a sensitivity analysis was carried out, which also explored the effect of excluding one article at a time on the overall risk calculation. Statistical significance was set at $P < 0.05$.

Results

Outcomes of the Search

The initial searches revealed a total of 37,793 papers within the Embase, PubMed, and Cochrane databases, comprising 7,359 from PubMed, 6,620 from Embase, and 23,814 from Cochrane. Following the removal of 6721 duplicate publications, 31,072 papers were evaluated through their titles and abstracts, leading to the marginalization of 25,480 distinctly irrelevant works, and 5592 articles underwent full-text evaluation. In the full-text screening process, 151 reviews or conference proceedings, 1799 animal experimental studies, 1816 duplicate cases, 1125 review articles, 565 case reports, and 74 correspondences were excluded. Finally, 62 articles were included for full-text reading and data extraction. Upon thorough examination of the complete text, 31 investigations that failed to satisfy the eligibility standards or lacked adequate data were subsequently excluded, and 31 studies were finally included for quantitative synthesis and meta-analysis (*Figure 1*). 12,227 people with diabetes were included in all included studies. Among them, 5256 males and 5887 females were explicitly reported, and the rest of the studies did not report gender composition. The included studies were geographically wide, including China, India, the Middle East, Europe, and the United States (*Table 1*) 14–44.

Quality evaluation

The NOS score of the investigations included in this analysis varied between 5 and 7, suggesting that the general standard of these investigations was considered satisfactory.

The meta-analysis of serum 25(OH) D₃ level

This review synthesized four independent studies to examine the serum 25(OH)D₃ levels in persons experiencing DR. Given the clear variability in the results ($I^2=99.5%$, $P=0.000$), a random-effects model was utilized for the combined evaluation. The mean of 25(OH)D₃ in the four studies was 10.68 ng/mL (95% CI: 5.94–15.41) after pooling. The particular outcomes are illustrated in *Figure 2*. To gauge the impact of particular investigations on the outcomes of the systematic review, we further performed sensitivity analyses. The consistency of the comprehensive meta-analysis findings can be assessed by

sequentially excluding individual studies and comparing the alterations in pooled effects and their confidence intervals before and after each exclusion. Sensitivity analysis showed that there were no obvious changes in the pooled mean value as a result of the exclusion of any other single study.

The meta-analysis of serum 1,25(OH)₂D₃

This study incorporated five independent investigations, and *Figure 3* illustrates the outcomes of a random-effects meta-analysis of serum 1,25(OH)₂D₃ amounts in persons experiencing DR. The results exhibited considerable variability ($I^2=98.8%$, $P<0.001$). The combined 1,25(OH)₂D₃ mean of the five studies was 31.14 pg/mL (95% CI: 24.35–37.94). The specific results are shown in *Figure 3*. We undertook sensitivity evaluations to assess the influence of particular investigations on the aggregate effect size of 1,25(OH)₂D₃ levels. The robustness of the results was investigated by excluding the included studies one by one and observing changes in the meta-analysis, projections of effect size. Sensitivity analysis showed that there were no obvious changes in the pooled mean value as a result of the exclusion of any other single study.

The meta-analysis of Serum Total 25(OH)D Levels

This meta-analysis included twenty-eight studies on serum total 25(OH)D amounts in those diagnosed with T2DM, conducted between 2012 to 2024. The mean serum Total 25(OH)D concentrations in individuals diagnosed with T2DM was 16.04 ng/mL (95% CI: 15.13–16.96 ng/mL) employing a random-effects model evaluation. The average total 25(OH)D amounts observed in the investigations varied between 6.49 ng/mL and 26.46 ng/mL. The investigations exhibited considerable variability ($I^2=98.8%$, $P<0.001$). Notably, most studies reported mean concentrations of 25(OH)D under 20 ng/mL indicate that deficiencies in vitamin D could be prevalent among individuals with T2DM. This discovery holds significant relevance for the medical management of diabetes. The specific results are depicted in *Figure 4*. We carried out sensitivity evaluations to gauge the influence of particular investigations on the overall effect size of Total 25(OH)D levels. The 28 included studies were sequentially excluded and alterations in the effect size estimates from the meta-analysis were noted to assess the reliability of the findings. Sensitivity analysis showed that there were no obvious changes in the pooled mean value as a result of the exclusion of any other single study.

Table 1 Characteristics of observational studies included in this meta-analysis.

Study	Country/ Region	Year	Subjects (male/female)	Mean age	Type of disease	Total 25 (OH)D	1,25(OH) ₂ D ₃		25(OH)D ₃	
Ahmed et al. (14)	Qatar	2021	460 (227/233)	55.2±15.0	T2DM	Total 25 (OH)D	26.46±15.30 ng/dL			
Ahmed et al. (15)	Qatar	2020	274 (110/164)	55.2±9.9	T2DM					25(OH)D ₃
Almoosa et al. (16)	Bahrain	2019	300 (NR)	NR	T2DM	Total 25 (OH)D	12.6±7.4 ng/mL			
Bajaj et al. (17)	India	2014	158 (95/63)	52.85±8.26	T2DM	Total 25 (OH)D	19.25±7.86 ng/mL			
Butler et al. (18)	Qatar	2020	460 (NR)	56.0±10.4	T2DM	Total 25 (OH)D	6.49±7.53 ng/mL	1,25(OH) ₂ D ₃	0.025±0.023 ng/mL	25(OH)D ₃
Castillo-Oti et al. (19)	Spanish	2022	385 (213/172)	69.89±9.86	T2DM	Total 25 (OH)D	15.50±10.80 ng/mL	1,25(OH) ₂ D ₃	24.50±13.00 pg/mL	
Choudhary et al. (20)	India	2024	119 (67/52)	49.5±11	T2DM	Total 25 (OH)D	16.3±9.2 ng/mL	1,25(OH) ₂ D ₃	42.5±15.6 pg/mL	
Hassan et al. (21)	Saudi Arabia	2023	252 (114/138)	18–87	T2DM	Total 25 (OH)D	15.50±7.60 ng/mL	1,25(OH) ₂ D ₃	37.40±9.20 pg/mL	
He et al. (22)	China	2014	1520 (773/747)	59.03±11.67	T2DM	Total 25 (OH)D	15.36±4.81 ng/mL			
Jung et al. (23)	South Korea	2016	257 (111/146)	58.8±12.1	T2DM	Total 25 (OH)D	14.4±8.6 ng/mL			
Wan et al. (24)	China	2019	4767 (2050/2717)	67±9	T2DM	Total 25 (OH)D	16.34±5.67 ng/mL			
Lopes et al. (25)	Portugal	2020	182 (86/96)	43±14	T1DM	Total 25 (OH)D	20.3±10.7 ng/mL			
Nadri et al. (26)	India	2021	88 (52/36)	53.18	T2DM	Total 25 (OH)D	14.07±1.21 ng/mL			
Nadri et al. (27)	India	2019	72 (NR)	53	T2DM	Total 25 (OH)D	14.10±1.20 ng/mL			
Navaei et al. (28)	Iran	2023	278 (93/185)	62.2±4.46	T2DM	Total 25 (OH)D	25.4±12.6 ng/mL			
Payne et al. (29)	United States	2012	42 (21/21)	59.8±12.0	T2DM	Total 25 (OH)D	21.1±10.5 ng/mL			
Reddy et al. (30)	India	2015	82 (48/34)	57.5±9.3	T2DM	Total 25 (OH)D	16.9±7.2 ng/mL			
Reheem et al. (31)	Egypt	2013	200 (96/104)	69±2.6	T2DM	Total 25 (OH)D	31.6±12.01 nmol/L	1,25(OH) ₂ D ₃	34.1±17.2 pmol/L	
Saxena et al. (32)	India	2019	66 (NR)	40-65	T2DM	Total 25 (OH)D	14.07±1.21 ng/mL			
Senyigit et al. (33)	Turkey	2019	163 (NR)	56.69±7.94	T2DM					25(OH)D ₃
Seyyar et al. (34)	Turkey	2022	165 (73/92)	62.5±6.8	T2DM	Total 25 (OH)D	11.77±7.86 ng/mL			
Seyyar et al. (35)	Turkey	2022	178 (87/91)	61.5±11.5	T2DM	Total 25 (OH)D	10.8±4.47 ng/mL			
Shimo et al. (36)	Japan	2014	75 (28/47)	28.5±5.5	T1DM					25(OH)D ₃
Usalp et al. (37)	Turkey	2023	40 (15/25)	58.9±7.8	T2DM	Total 25 (OH)D	12.8±4.2 ng/mL			
Usluogullari et al. (38)	Turkey	2015	557 (296/261)	55.2±10.9	T2DM	Total 25 (OH)D	19.1±8.1 ng/mL			
Vaghela et al. (39)	India	2022	98 (57/43)	58.9±13.6	T2DM	Total 25 (OH)D	14.4±5.9 ng/mL			
Verma et al. (40)	India	2024	98 (52/46)	57.6±10.4	T2DM	Total 25 (OH)D	25.91±7.83 ng/mL			
Yuan et al. (41)	China	2019	273 (143/130)	58.10±10.83	T2DM	Total 25 (OH)D	13.14±8.71 ng/mL			
Zahedi et al. (42)	Iran	2024	201 (134/68)	60.0±8.1	T2DM	Total 25 (OH)D	14.46±7.88 ng/mL			
Zhao et al. (43)	China	2021	235 (120/89)	61±6.8	T2DM	Total 25 (OH)D	16.38±9.16 ng/mL			
Zhuang et al. (44)	China	2024	182 (95/87)	51.1±7.6	T2DM	Total 25 (OH)D	17.34±4.40 ng/mL			




Figure 1 Document retrieval flow chart.




Figure 2 Forest map of serum 25(OH)D₃ levels in diabetic retinopathy patients.

Publication bias assessment

We assessed the likelihood of publication bias in the investigations that were included through plotting funnel plots for serum 25(OH)D₃, 1,25(OH)₂D₃, and Total 25(OH)D (Figure 5). Visually, the funnel plot's left and right sides are nearly symmetrical. In order to quantitatively evaluate the extent of bias in publica-

tions, we conducted Egger's linear regression test for each of the three indicators. The results showed that the P values of Egger's test for 25(OH)D₃, 1,25(OH)₂D₃ and Total 25(OH)D were 0.466, 0.908 and 0.727, respectively, which were greater than the significance threshold of 0.05. This indicates that no obvious publication bias was observed in this meta-analysis.




Figure 3 Forest map of serum 1,25(OH)₂D₃ levels in diabetic retinopathy patients.




Figure 4 Forest map of serum Total 25(OH)D levels in diabetic retinopathy patients.




Figure 5 Serum 25(OH)D₃, 1,25(OH)₂D₃ and Total 25(OH)D funnel plot.

Discussion

This study employed a meta-analysis and systematic review to measure the evidence regarding vitamin D levels among individuals suffering from DR. The findings indicated that serum amounts of 25(OH)D₃, 1,25(OH)₂D₃, and Total 25(OH)D in persons with DR were lower, implying an intriguing link among insufficient amounts of vitamin D and the initiation and advancement of DR, which aligns with some previous studies (5, 6, 10). The findings were broadly consistent, but the results were more convincing as more high-quality studies were included in this study.

DR is a prevalent microvascular problem associated with diabetes, and its development is influenced by a range of factors (45, 46). Vitamin D plays part in the pathogenic course of DR through various processes. First, vitamin D plays a role in inhibiting the inflammatory response and reducing the production

of cytokines that are pro-inflammatory, thus reducing inflammatory damage to retinal micro vessels. Secondly, vitamin D has an antioxidant effect, which can remove excess reactive oxygen species (ROS) and safeguard retinal pigment epithelial cells and ganglion cells (47). Thirdly, vitamin D can enhance the synthesis of anti-vascular endothelial growth factor and suppress pathological neovascularization (48). Moreover, vitamin D enhances the function of vascular smooth muscle cells and mitigates microcirculation disorders via management of calcium ion influx (49). Considering the various protective benefits associated with vitamin D, correcting vitamin D insufficiency is expected to impede the advancement of DR. The results of this study showed that vitamin levels in patients with DR were significantly lower than those in patients with non-diabetic retinopathy. This suggests that retinopathy in diabetic patients may be closely related to vitamin D deficiency and that our findings provide strong evidence for guiding the clinical man-

agement of diabetic patients, thereby helping to reduce retinopathy.

The findings of the systematic review conducted in this study indicated a notable level of heterogeneity among the investigations that were included, with I^2 values exceeding 98%. Heterogeneity can stem from a variety of factors. First, participants in the studies included exhibited variations in baseline characteristics, including age, gender, ethnicity, duration of diabetes, and comorbidities. These confounding factors may influence vitamin D quantities and bring about discrepancies in effect sizes across studies (50). Secondly, the diagnostic criteria for DR in each study were different, some studies used fundus photography, some used fluorescein fundus angiography, and some studies did not provide a clear basis for diagnosis, and the heterogeneity of DR diagnosis directly led to differences in patient characteristics (51). Thirdly, significant differences exist in the detection methods for serum 25(OH)D₃, 1,25(OH)₂D₃, and Total 25(OH)D. The techniques encompass liquid chromatography-mass spectrometry, radioimmunoassay, and chemiluminescence immunoassay (CLIA). Performance indicators such as sensitivity, specificity, and detection limits vary considerably among these methods, potentially resulting in systematic errors in vitamin D level assessments (52). In addition, the geographical distribution of the included studies is wide, covering different countries and places like China, India, the Middle East, Europe, and United States, and variations in race, eating habits, and sunshine hours could influence the manufacturing and breakdown of vitamin D in the human body (53). Subgroup analysis indicated that levels of 25(OH)D₃ and Total 25(OH)D were elevated in European and American populations compared to the Asian population ($P < 0.05$), highlighting the significance of ethnic and regional differences as sources of heterogeneity². Finally, the sample sizes of the investigations included in the analysis exhibited considerable variability, with smaller studies demonstrating a greater propensity to yield extreme outcomes, resulting in a deviation from the true effect size estimates (54). Considering the above factors, the high heterogeneity reduces the comparability and robustness of the results of this study to a certain extent.

This study has certain limitations. The studies included were all single arm, making it challenging to ascertain the link of causation between the quantities of vitamin D and DR owing to the absence of appropriate control groups. Reduced quantities of vitamin D may be a consequence of DR, or both could be related to other factors. Prospective cohort studies can help to further clarify the relationship between the two. Second, funnel plots suggest that there may be publication bias, and negative studies with small sample sizes may be difficult to publish, resulting in results that deviate from the true effect. Finally, the

quality of individual studies varied and may be affected by selection bias, information bias, and confounding bias. We assessed only the overall quality of the studies during the inclusion process, lacking a more detailed assessment of the danger of bias. The limitations may have influenced the dependability of the investigation's findings. From the above limitations, future research directions include: (1) conducting cohort studies with large samples and extensive follow-up to assess the correlation between low levels of vitamin D and the risk of DR; (2) to explore the differences of vitamin D in patients with different stages and types of DR, along with the dose-response connection between vitamin D and the seriousness of DR; (3) carry out clinical intervention investigations into vitamin D supplemental intake to evaluate the impact of vitamin D therapy on the advancement of DR; Comprehensive examination of the molecular mechanism connected to low-level vitamin D promoting the existence and growth of DR. These studies elucidate the significance of vitamin D in both the avoidance and control of DR and provide innovative ideas for the development of intervention strategies.

Conclusion

In conclusion, the study demonstrates that a lack of vitamin D is often found in individuals with DR, implying that vitamin D may be involved in DR development. Given the prominent level of heterogeneity and potential for publication bias, more high-quality prospective investigations are required to substantiate this conclusion. In future studies, it is essential to assess the link between reduced amounts of vitamin D and prognosis of DR, along with exploring the potential clinical utilization of vitamin D in both the avoidance and control of DR.

Funding

This work was supported by the Optimization of Integrated Traditional Chinese and Western Medicine Diagnosis and Treatment Plan for Retinal Artery Occlusion and Construction of Intelligent Auxiliary Diagnosis and Treatment System (NO.23377712D); The High-level Key Discipline Construction Project of the National Administration of Traditional Chinese Medicine (zyzdxk-2023017); Xingtai City Innovation Capacity Enhancement Program Project (No. 2023ZZ106); Innovation Support Project for Post-graduates in Hebei Province (XCXZZBS2025026).

Conflict of interest statement

All the authors declare that they have no conflict of interest in this work.

References

- Wong TY, Cheung CM, Larsen M, Sharma S, Simo R. Diabetic retinopathy. *Nat Rev Dis Primers* 2016; 2: 16012.
- Yau JW, Rogers SL, Kawasaki R, Lamoureux EL, Kowalski JW, Bek T, et al. Global prevalence and major risk factors of diabetic retinopathy. *Diabetes Care* 2012; 35(3): 556–64.
- Cheung N, Mitchell P, Wong TY. Diabetic retinopathy. *Lancet* 2010; 376(9735): 124–36.
- Holick MF. Vitamin D deficiency. *New Engl J Med* 2007; 357(3): 266–81.
- Luo BA, Gao F, Qin LL. The Association between Vitamin D Deficiency and Diabetic Retinopathy in Type 2 Diabetes: A Meta-Analysis of Observational Studies. *Nutrients* 2017; 9(3):
- Bikle DD. Vitamin D metabolism, mechanism of action, and clinical applications. *Chem Biol* 2014; 21(3): 319–29.
- Zittermann A, Gummert JF. Nonclassical vitamin D action. *Nutrients* 2010; 2(4): 408–25.
- Payne JF, Ray R, Watson DG, Delille C, Rimler E, Cleveland J, et al. Vitamin D insufficiency in diabetic retinopathy. *Endocr Pract* 2012; 18(2): 185–93.
- Aksoy H, Akcay F, Kurtul N, Baykal O, Avci B. Serum 1,25 dihydroxy vitamin D (1,25(OH)2D3), 25 hydroxy vitamin D (25(OH)D) and parathormone levels in diabetic retinopathy. *Clin Biochem* 2000; 33(1): 47–51.
- Bajaj S, Singh RP, Dwivedi NC, Singh K, Gupta A, Mathur M. Vitamin D levels and microvascular complications in type 2 diabetes. *Indian J Endocrinol Metab* 2014; 18(4): 537–41.
- Bonakdaran S, Shoeibi N. Is there any correlation between vitamin D insufficiency and diabetic retinopathy? *Int J Ophthalmol-Chi* 2015; 8(2): 326–31.
- Wilkinson CP, Ferris FR, Klein RE, Lee PP, Agardh CD, Davis M, et al. Proposed international clinical diabetic retinopathy and diabetic macular edema disease severity scales. *Ophthalmology* 2003; 110(9): 1677–82.
- Margulis AV, Pladevall M, Riera-Guardia N, Varas-Lorenzo C, Hazell L, Berkman ND, et al. Quality assessment of observational studies in a drug-safety systematic review, comparison of two tools: the Newcastle-Ottawa Scale and the RTI item bank. *Clin Epidemiol* 2014; 6: 359–68.
- Ahmed L, Butler AE, Dargham SR, Latif A, Ahmed EA, Hassan A, et al. Relationship between total vitamin D metabolites and complications in patients with type 2 diabetes. *Biomed Rep* 2021; 14(1): 18.
- Ahmed L, Butler AE, Dargham SR, Latif A, Robay A, Chidiac OM, et al. Association of vitamin D(2) and D(3) with type 2 diabetes complications. *Bmc Endocr Disord* 2020; 20(1): 65.
- Abdulla Almoosa, Seemantini Ayachit, Abdulaziz Aldoseri, Wael Wagih. Incidence of Vitamin D Deficiency in Patients with Type II Diabetes Mellitus and its Relation to the Severity of Retinopathy. *Bahrain Medical Bulletin* 2019; 41: 4
- Bajaj S, Singh RP, Dwivedi NC, Singh K, Gupta A, Mathur M. Vitamin D levels and microvascular complications in type 2 diabetes. *Indian J Endocrinol Metab* 2014; 18(4): 537–41.
- Butler AE, Dargham SR, Latif A, Mokhtar HR, Robay A, Chidiac OM, et al. Association of vitamin D(3) and its metabolites in patients with and without type 2 diabetes and their relationship to diabetes complications. *Ther Adv Chronic Dis* 2020; 11: 1754215935.
- Castillo-Oti JM, Galvan-Manso AI, Callejas-Herrero MR, Vara-Gonzalez LA, Salas-Herrera F, Munoz-Cacho P. Vitamin D Deficiency Is Significantly Associated with Retinopathy in Type 2 Diabetes Mellitus: A Case-Control Study. *Nutrients* 2021; 14(1):
- Shrikant Choudhary, Shubham Gupta, Anil kumar Panwar, Animesh Gaur, Ramkishan Jat. Association of Vitamin D Level With Micro Vascular Complications of Type 2 Diabetes Mellitus. *Journal of Cardiovascular Disease Research* 2024; 15: 1710–7.
- Tahani Hassan, Nadine Alhumaidan, Basher Yousef, Mariam Alsalthi, Abdulrahman Albulayhi, Sultan Al Harbi. The Association between Vitamin D Status and Diabetic Complications in Type 2 Diabetic Patients. *Biomedical & Pharmacology Journal* 2023; 16: 2469–75.
- He R, Shen J, Liu F, Zeng H, Li L, Yu H, et al. Vitamin D deficiency increases the risk of retinopathy in Chinese patients with type 2 diabetes. *Diabetic Med* 2014; 31(12): 1657–64.
- Jung CH, Kim KJ, Kim BY, Kim CH, Kang SK, Mok JO. Relationship between vitamin D status and vascular complications in patients with type 2 diabetes mellitus. *Nutr Res* 2016; 36(2): 117–24.
- Wan H, Wang Y, Zhang K, Chen Y, Fang S, Zhang W, et al. Associations Between Vitamin D And Microvascular Complications In Middle-Aged And Elderly Diabetic Patients. *Endocr Pract* 2019; 25(8): 809–16.
- Lopes M, Laiginhas R, Madeira C, Neves JS, Barbosa M, Rosas V, et al. Association between Serum Vitamin D and Diabetic Retinopathy in Portuguese Patients with Type 1 Diabetes. *Acta Medica Port* 2020; 33(7–8): 459–65.
- He J, Kang J, Mei Z, Zhou X, Yin Y, Zhu C. Gender-specific association between circulating 25-hydroxyvitamin D levels and diabetic retinopathy in patients with type 2 diabetes mellitus. *Front Endocrinol* 2025; 16: 1541961.
- Nadri G, Saxena S, Mahdi AA, Kaur A, Ahmad MK, Garg P, et al. Serum vitamin D is a biomolecular biomarker for proliferative diabetic retinopathy. *Int J Retina Vitre* 2019; 5: 31.
- Navaei S, Nazemi S, Emamian MH, Hashemi H, Fotouhi A. Vitamin D deficiency and diabetic retinopathy risk. *J Fr Ophthalmol* 2023; 46(7): 737–41.
- Payne JF, Ray R, Watson DG, Delille C, Rimler E, Cleveland J, et al. Vitamin D insufficiency in diabetic retinopathy. *Endocr Pract* 2012; 18(2): 185–93.
- Reddy GB, Sivaprasad M, Shalini T, Satyanarayana A, Seshacharyulu M, Balakrishna N, et al. Plasma vitamin D status in patients with type 2 diabetes with and without retinopathy. *Nutrition* 2015; 31(7–8): 959–63.

31. Rania Naguib Abdel Mouteleb Abdel Reheem, Maaly Abdel Halim Mohamed Abdel Fattah b. Serum vitamin D and parathormone (PTH) concentrations as predictors of the development and severity of diabetic retinopathy. *Alexandria Journal of Medicine* 2013; 49: 119–23.
32. Saxena S, Nadri G, Kaur A, Mahdi A, Ahmad K, Garg P. Low serum vitamin D levels correlate with disorganization of retinal inner layers, ellipsoid zone disruption and retinal pigment epithelium alterations in diabetic retinopathy. *Invest Ophth Vis Sci* 2019; 60: 5321.
33. Senyigit A. The association between 25-hydroxy vitamin D deficiency and diabetic complications in patients with type 2 diabetes mellitus. *Diabetes Metab Syndr* 2019; 13(2): 1381–6.
34. Seyyar SA, Tiskaoglu NS, Onder TE, Mercanli M, Dogan L. Is serum vitamin D associated with diabetic retinopathy and its severity or with diabetes itself? *Clin Exp Optom* 2023; 106(6): 612–8.
35. Seyyar SA, Tokuc EO, Tiskaoglu NS, Karabas VL, Gungor K. Do serum vitamin D levels correlate with Macular Edema or with Diabetic Retinopathy? *Eur J Ophthalmol* 2022; 32(6): 3592–8.
36. Shimo N, Yasuda T, Kaneto H, Katakami N, Kuroda A, Sakamoto F, et al. Vitamin D deficiency is significantly associated with retinopathy in young Japanese type 1 diabetic patients. *Diabetes Res Clin Pr* 2014; 106(2): e41–3.
37. Zuhaf Onur Usalp. Is there a relationship between diabetic retinopathy and vitamin D? *Ann Clin Anal Med* 2023; 14: 332–5.
38. Usluogullari CA, Balkan F, Caner S, Ucler R, Kaya C, Ersoy R, et al. The relationship between microvascular complications and vitamin D deficiency in type 2 diabetes mellitus. *Bmc Endocr Disord* 2015; 15: 33.
39. Vikas K Vaghela, Krushna Kishore Padala, Keyur Madhu. Status and correlation of 25-hydroxy Vitamin D in diabetes mellitus Type II patients with and without retinopathy. *National Journal of Physiology, Pharmacy and Pharmacology* 2022; 12: 10.
40. Chhaya Verma. Inverse Relation Of 25-Oh Vitamin D Level With Severity Of Diabetic Retinopathy And Hba1c In Type 2 Diabetes Patients: Supplementation May Halt Severity. *Asian J Pharm Clin Res* 2024; 17: 112–5.
41. Yuan J, Zhou JB, Zhao W, Zhang RH, Cai YH, Shu LP, et al. Could Vitamin D be Associated with Proliferative Diabetic Retinopathy? Evidence from Pooling Studies. *Horm Metab Res* 2019; 51(11): 729–34.
42. Zahedi M, Motahari MM, Fakhri F, Aphshari NM, Poursharif S, Jahed R, et al. Is vitamin D deficiency associated with retinopathy in type 2 diabetes mellitus? A case-control study. *Clin Nutr Espen* 2024; 59: 158–61.
43. Zhao WJ, Xia XY, Yin J. Relationship of serum vitamin D levels with diabetic microvascular complications in patients with type 2 diabetes mellitus. *Chinese Med J Peking* 2021; 134(7): 814–20.
44. Zhuang Y, Zhuang Z, Cai Q, Hu X, Huang H. Serum vitamin D is substantially reduced and predicts flares in diabetic retinopathy patients. *J Diabetes Invest* 2024; 15(7): 867–73.
45. Dai L, Sheng B, Chen T, Wu Q, Liu R, Cai C, et al. A deep learning system for predicting time to progression of diabetic retinopathy. *Nat Med* 2024; 30(2): 584–94.
46. Pushparani DS, Varalakshmi J, Roobini K, Hamshapriya P, Livitha A. Diabetic Retinopathy-A Review. *Curr Diabetes Rev* 2025; 21(7): 43–55.
47. Tohari AM, Alhasani RH, Biswas L, Patnaik SR, Reilly J, Zeng Z, et al. Vitamin D Attenuates Oxidative Damage and Inflammation in Retinal Pigment Epithelial Cells. *Antioxidants-Basel* 2019; 8(9):
48. Albert DM, Scheef EA, Wang S, Mehraein F, Darjatmoko SR, Sorenson CM, et al. Calcitriol is a potent inhibitor of retinal neovascularization. *Invest Ophth Vis Sci* 2007; 48(5): 2327–34.
49. Ren Z, Li W, Zhao Q, Ma L, Zhu J. The impact of 1,25-dihydroxy vitamin D₃ on the expressions of vascular endothelial growth factor and transforming growth factor-beta(1) in the retinas of rats with diabetes. *Diabetes Res Clin Pr* 2012; 98(3): 474–80.
50. Azizieh F, Alyahya KO, Raghupathy R. Association between levels of vitamin D and inflammatory markers in healthy women. *J Inflamm Res* 2016; 9: 51–7.
51. Zhang J, Upala S, Sanguankeo A. Relationship between vitamin D deficiency and diabetic retinopathy: a meta-analysis. *Can J Ophthalmol* 2017; 52(2): 219–24.
52. Wallace AM, Gibson S, de la Hunty A, Lamberg-Allardt C, Ashwell M. Measurement of 25-hydroxyvitamin D in the clinical laboratory: current procedures, performance characteristics and limitations. *Steroids* 2010; 75(7): 477–88.
53. Hilger J, Friedel A, Herr R, Rausch T, Roos F, Wahl DA, et al. A systematic review of vitamin D status in populations worldwide. *Brit J Nutr* 2014; 111(1): 23–45.
54. von Hippel PT. The heterogeneity statistic I(2) can be biased in small meta-analyses. *Bmc Med Res Methodol* 2015; 15: 35.

Received: July 25, 2025

Accepted: September 09, 2025